

- cer. *World J Surg* 1998;22:223-7; discussion 227-8.
- (37) Adler LP, Crowe JP, al-Kaisi NK, Sunshine JL. Evaluation of breast masses and axillary lymph nodes with [F-18] 2-deoxy-2-fluoro-D-glucose PET. *Radiology* 1993;187:743-50.
- (38) Wahl RL, Henry CA, Ethier SP. Serum glucose: effects on tumor and normal tissue accumulation of 2-[F-18]-fluoro-2-deoxy-D-glucose in rodents with mammary carcinoma. *Radiology* 1992;183:643-7.
- (39) Crippa F, Gavazzi C, Bozzetti F, Chiesa C, Pascali C, Bogni A, et al. The influence of blood glucose levels on [<sup>18</sup>F]fluorodeoxyglucose (FDG) uptake in cancer: a PET study in liver metastases from colorectal carcinomas. *Tumori* 1997;83:748-52.
- (40) van der Wall E. The sentinel node in breast cancer: implication for adjuvant treatment? *Eur J Nucl Med* 1999;26(4 Suppl):S17-9.
- (41) Oshida M, Uno K, Suzuki M, Nagashima T, Hashimoto H, Yagata H, et al. Predicting the prognoses of breast carcinoma patients with positron emission tomography using 2-deoxy-2-fluoro[<sup>18</sup>F]-D-glucose. *Cancer* 1998;82:2227-34.

## NOTES

Supported in part by the Progetto Finalizzato Ministero Sanità and the Associazione Italiana Ricerca sul Cancro.

We are deeply indebted to Dr. Marco Sandri (Center for Scientific Calculus, University of Verona, Italy) for mathematical help.

Manuscript received July 12, 2000; revised January 30, 2001; accepted February 7, 2001.